

Dermatology Associates

Mohs Micrographic Surgery

Johnson City
2885 Boones Creek Road
423-928-9014

Kingsport
2300 W Stone Drive
423-246-4961 • 1-800-445-7274

Bristol
3183 W State Street
423-764-7131

In preparation for your upcoming appointment for Mohs surgery, please review the enclosed information packet. This information will give you some insight into Mohs micrographic surgery, answer some frequently asked questions, and provide directions to our office. Our hope is that this information will answer many of your questions and relieve potential anxiety as your appointment approaches.

Important information:

- Questions concerning your upcoming surgery can be addressed by our Mohs Support Specialist:

Johnson City
423-928-8088

Kingsport
423-230-3145

Bristol
423-764-7131 ext 5094

- Continue taking all of your regular medications including aspirin or blood thinning medications if these have been prescribed by your physician.
- If you take antibiotics prior to dental or other surgical procedures, or have an artificial heart valve, please mention this once you arrive for your appointment. We may give antibiotics prior to the surgery if necessary.
- Anticipate quite a few stitches. In order to have a superior final cosmetic result it often requires multiple tiny sutures that usually extend outside the area where the cancer is located. Expect bruising and swelling for several weeks after the surgery.
- Plan on being in our office a good portion of the day.

On the day of your surgery:

- **Bring a driver for your return home**
- **Take all of your normal medications as prescribed**
- **Do not apply makeup, creams, lotions, etc. near the surgery site**
- **Eat a healthy breakfast and bring snacks / lunch**

*****The Mohs surgery section of our website has an excellent 4 min video explaining Mohs surgery – www.tricitiesderm.com**

Have confidence in knowing you will be seeing a Fellowship Trained Mohs Surgeon – most Mohs surgeons are NOT fellowship trained. We look forward to making sure your experience with Mohs surgery is a positive one and appreciate the opportunity to serve you.

Mohs Micrographic Surgery

Mohs surgery, also called Mohs micrographic surgery, is a specialized technique for removing skin cancer. Named for Dr. Frederic Mohs, who developed the technique, Mohs surgery is the treatment of choice for many skin cancers. It offers the following advantages:

- Highest cure rate. The tissue is prepared and examined in our lab which allows for complete microscopic evaluation of 100% of the deep and lateral margins. This allows mapping of the cancer to precisely trace out and remove all the roots.
- Best cosmetic result. Because there is no need to take a large margin of healthy skin in Mohs surgery, the size of the wound left after tumor removal is minimized. Keeping the defect small optimizes the surgical repair, and produces a superior cosmetic result.

Mohs surgery is an outpatient procedure that is performed under local anesthesia; therefore the risks associated with prolonged general anesthesia (being “put to sleep”) are avoided. Before the surgery begins, a small needle is used to place the numbing medicine in the area surrounding the skin cancer. After this first step, the remainder of the procedure is relatively pain free. The surgeon first removes the obvious skin cancer that can be seen with the unaided eye. Next, a very thin saucer-shaped layer of normal appearing skin is removed taking special care to map the tissue so that the surgeon is able to determine the corresponding margins on the patient. The tissue is then processed by the laboratory located in our office – a process that takes about an hour. Microscopic slides are prepared by a technologist and examined by the surgeon to determine if the cancer is persistent.

If microscopic examination reveals remaining tumor, a map is drawn indicating the precise location. You are then brought back to the surgery suite, and additional anesthetic is injected to reinforce the initial injection. In most cases, the area is still numb and you will feel little to no discomfort.

Another layer of tissue is removed – but only where the map indicated residual cancer. The healthy tissue is left alone and only the cancer-containing tissue is excised. This tissue is again brought to the laboratory and the process is repeated until all evident cancer is removed.

Because the number of stages involved with each case is unpredictable, it is impossible to know how long the surgery will take. For this reason, it is important for patients to plan on spending most of the day with us. Do not be discouraged if your cancer is not removed in one step. Remember, we are tracing the roots of the tumor in a stepwise fashion. This must be done in small layers. Bring a good book and whatever else will help keep you comfortable while you are with us.

Once all the cancer has been removed, we will then discuss the options to reconstruct the resulting defect or open wound. This usually requires stitches.

Frequently Asked Questions

What does "Mohs" stand for?

Dr. Frederic Mohs developed this technique about 60 years ago. The procedure has been modified and refined over the years. Practitioners of the technique have kept Dr. Mohs' name in respect for his contribution. Mohs surgery has other names including Mohs chemosurgery, Mohs microscopically controlled surgery, and Mohs micrographic surgery.

Are there different types of skin cancer?

Yes. The most common type of skin cancer is basal cell carcinoma. Often these skin cancers are very subtle, and appear like a small pimple that will not heal and often bleed. The second most common skin cancer is squamous cell carcinoma. This type usually appears as a thickened scaly plaque or nodule. The third most common type is melanoma, which behaves differently than either basal cell carcinoma or squamous cell carcinoma. Melanoma most commonly appears as an abnormal mole.

Are skin cancers life-threatening?

Fortunately, the two most common types of skin cancer, basal cell carcinoma and squamous cell carcinoma, are rarely life threatening. As a general rule, basal cell carcinoma does not spread to the glands or other parts of the body. Compared to basal cell carcinoma, squamous cell carcinoma has an increased risk to spread to other parts of the body. Fortunately, distant spread is uncommon when the cancer is treated early. However, both types of tumors will continue to grow locally and destroy normal tissue. The third most common type of skin cancer, melanoma, can be life threatening if treated late. Basal cell skin cancers and squamous cell cancers never "turn into" melanoma.

How large of a scar will I have from the surgery?

The size of the scar depends on the size of the tumor. It is often difficult to predict the size of the tumor prior to surgery.

Will I have stitches following the surgery?

Most patients will have stitches. Often patients are surprised at the number of stitches required or the length of the incision. Please keep in mind that our goal is the best long term cosmetic result. This may initially require extensive suturing. There are four main ways your surgical wound may be handled:

1. Direct closure of wound with stitches. This is the most common method.
2. Skin flap: In some instances, skin that is located near the wound is moved in to fill the defect left from surgery.
3. Skin graft: In certain situations it is appropriate to take skin from a site distant from the wound (usually in front or behind the ear, or from the neck) and use it to cover the surgical defect.
4. Let wound heal in by itself. The body has an excellent capacity to heal open wounds. This healing period is approximately three to six weeks depending on the size of the wound. It requires regular wound care.

In addition to wound size and location, the surgeon considers other factors in determining how your wound will be handled. This will be fully discussed with you on the day of surgery.

Will I be put to sleep for the surgery?

No. The surgery is well tolerated with local anesthesia. Because the surgery may be time-consuming, the risk of prolonged general anesthesia is thus avoided.

How long will the surgery last?

The length of surgery depends on the extent of the tumor. Often surgery lasts half a day or longer. Much of the time is spent waiting for tissue to be processed. Bring reading materials, needlework, etc., with you to help pass the time. Also, you may want to bring a snack or lunch with you on the day of surgery.

What if I live far away from the area?

If your travel distance is great you might want to spend the night before surgery in the local area. There are a few moderately priced hotels nearby. A list of these is available upon request.

Should I bring someone with me?

It is often helpful to bring someone with you on the day of surgery. Depending on the size and location of your tumor, driving is sometimes a challenge after surgery, and having a driver with you is a good idea. Additionally, if someone other than the patient will be performing post-op wound care, the nurse can give this person direct instructions, demonstrate wound care, and answer any questions about the surgery.

What should I wear?

You should wear comfortable clothing. You may want to bring a sweater, as our office is kept cool for the laboratory equipment that is needed for this procedure.

Should I eat breakfast before surgery?

Yes. Breakfast is recommended. The only exception would be if you are scheduled to have your reconstruction done by another physician after Mohs has been completed. In this case you will follow the instructions from his/her office.

Should I take my regular medications on the morning of the surgery?

Yes. Take your regular medications as they have been prescribed.

Are there any medications I should avoid prior to surgery?

If you take aspirin on a regular basis at the direction of your physician, you SHOULD CONTINUE to take it as prescribed. If you have decided yourself to take aspirin, NOT at the recommendation of a physician, it may be beneficial to stop your aspirin 10 days prior to surgery. If you are on any other blood thinners (Coumadin® or warfarin, Plavix®, Pradaxa®, etc) please continue taking them as prescribed.

Will my activity be limited after surgery?

Yes. You will need to keep your heart rate normal for 48 hours after the surgery. Physical activity, including sports, may be limited for the first week. If your work requires significant physical exertion, you may consider taking a few days off following the surgery. Avoid any long trips within the first few days following surgery in case you develop any complications. If you have any questions or special situations please do not hesitate to call our office.

Since I have taken antibiotics before other procedures in the past do I need to take them before Mohs surgery?

If you have taken antibiotics prior to dental or other surgical procedures, or have artificial heart valve, we may give a single dose of antibiotics prior to Mohs surgery. This will help prevent infection of the heart valve. Pacemakers, implanted defibrillators, mitral valve prolapse and cardiac stents do not generally require antibiotics. The antibiotic used for dental work is best for bacteria found in the mouth and is not as effective for skin surgery. Please inform our nurses upon arrival if you have been told to take antibiotics prior to a procedure.

What are the potential complications of surgery?

Bleeding and infection are the two primary complications. Both of these are uncommon, and can be treated if they do occur. We will discuss how to recognize and deal with these problems when you come for your surgery.

Will my insurance cover the cost of surgery?

Under most circumstances your carrier will pay for surgery. If you are in doubt about your particular coverage, you should check with your insurance representative prior to your appointment. If you have specific questions regarding insurance or billing matters, please contact our office at 423-246-4961.

Will my surgery qualify for FMLA?

In general, no. Only in extreme circumstances would a Mohs procedure qualify for FMLA.

Learn more about Mohs surgery from:

American College of Mohs Surgery

www.mohscollege.org/

www.skincancermohssurgery.org/

www.tricitiesderm.com

We look forward to making sure your experience with Mohs surgery is a positive one. Please do not hesitate to call with any questions or concerns.

Dermatology Associates
Mohs Micrographic Surgery

Chad J Thomas, MD - cthomas@tricitiesderm.com

Erin Reid, MD - ereid@tricitiesderm.com

Cory Trickett, DO - ctrickett@tricitiesderm.com

CJT 2/19